### Heavy-tailed distributions

# References

- [1] V.P. Chistyakov (1964). A theorem on sums of independent, positive random variables and its applications to branching processes. *Theory Probab. Appl.* **9**, 640–648.
- [2] P. Embrechts, C. Klüppelberg & T. Mikosch, T (1997). Modelling Extremal Events for Insurance and Finance. Springer-Verlag, Heidelberg.
- [3] C. Klüppelberg (1988). Subexponential distributions and integrated tails. J. Appl. Probab. 25, 132–141.
- [4] K. Sigman (1999). A primer on heavy-tailed distributions, QUESTA, 33, 261–275.

### Single-server delay results

# References

- S. Asmussen, H. Schmidli and V. Schmidt (1999). Tail Probabilities for Non-Standard Risk and Queueing Processes with Subexponential Jumps. Adv. Appl. Probab., 31.
- [2] J.W. Cohen (1973). Some results on regular variation for distributions in queueing and fluctuation theory. QUESTA, 33, 205–232.
- [3] P. R. Jelenković and A. A. Lazar (1998). Subexponential asymptotics of a Markovmodulated random walk with queueing applications. J. Appl. Probab., **35**, 325–347.
- [4] A. Pakes (1975). On the tails of waiting time distributions. J. Appl. Probab. 12, 555–564.
- [5] W.L. Smith (1972). On the tails of queueing time distributions. (Technical Report, MIMEO Series No. 830, Dept. Statistics, U. North Carolina, Chapel Hill.)

### Tandem queues and beyond

# References

- [1] O.J. Boxma and Q. Deng (2000). Asymptotic properties in the tandem queueing system with identical service times at both queues. *Mathematical Methods of Operations Research*, Volume **52**).
- [2] F. Baccelli, S. Schlegel and V. Schmidt (1999). Asymptotics of stochastic networks with subexponential service times. QUESTA, 33,205–232.
- [3] T. Huang, and K. Sigman (1999). Steady-state asymptotics for tandem, split-match and other feedforward queues with heavy-tailed service. *QUESTA*, **33**,233–260.

#### Queue length, busy periods and processor sharing

# References

- S. Asmussen, C. Kluppelberg and K. Sigman (1999). Sampling at subexponential times, with queueing applications. *Stoch. Proc Appls.* 79, 265–286.
- [2] S. Borst, O.J. Boxma and P. R. Jelenković (2002). Reduced-load equivalence and induced burstiness in GPS queues with long-tailed traffic flows. *QUESTA* (to appear)
- [3] P. R. Jelenković and P. Momcilovic, (2002). Large deviation analysis of subexponential waiting times in a processor sharing queue. *Math. Opns. Res.* (to appear)
- [4] A. de Meyer and J.L.T. Teugels (1980). On the asymptotic behaviour of the distributions of the busy period and service time in M/G/1. J. Appl. Probab. 17, 820–813.
- [5] R. Nunez-Queija (2000) Processor sharing models for integrated service networks. (Ph.D. dissertation, Eindhoven University of Technology)
- [6] A.P. Zwart and O.J. Boxma (2000) Sojourn time asymptotics in the M/G/1 processor sharing queue. Syst. Theory Appl. 35, 141–166.
- [7] A.P. Zwart (2001) Tail asymptotics for the busy period in the GI/GI/1 queue. Math. Opns. Res. 26, 485–493.

### Fluid queues

### References

- R. Agrawal, A.M. Makowski and P. Nain (1999). On a reduced load equivalence for fluid queues under subexponentiality. *QUESTA*, 33, 5–42.
- [2] P. R. Jelenković and P. Momcilovic, and B. Zwart (2001). Reduced load equivalence under subexponentiality.
- [3] P. R. Jelenković and P. Momcilovic, (2002). Asymptotic loss probability in a finite buffer fluid queue with heterogeneous heavy-tailed on-off processes. *Annals Appl. Probab.* (to appear)
- [4] P. R. Jelenković and P. Momcilovic, and B. Zwart (2001). Reduced load equivalence under subexponentiality.
- [5] S. Resnick and G. Samorodnitsky (1999). Activity periods of an infinite server queue and performance of certain heavy tailed fluid queues. *QUESTA*, **33**,43–72.
- [6] A.P. Zwart, S.C. Borst and M. Mandjes (2000) Exact asymptotics for fluid queues fed by multiple heavy-tailed on-off flows. (Technical Report, University of Eindhoven).

# Misc

# References

- J. Abate, G.L. Choudhury and W. Whitt(1995). Exponential approximations for tail probabilities of queues, I: waiting times. *Opns. Res.*, 43, 885–901.
- [2] B. von Bar (1975). Asymptotic ruin probabilities when exponential moments do not exist. Scand. Act. J. 6-10. A theorem on sums of independent, positive random variables and its applications to branching processes. Theory Probab. Appl. 9, 640–648.
- [3] M. Bramson (1994). Instability of FIFO queueing networks, Ann. Appl. Probab., 4, 414–431.
- [4] P. Embrechts & N. Veraverbeke (1982). Estimates for the probability of ruin with special emphasis on the possibility of large claims. *Insurance: Mathematics and Economics* 1, 55–72.
- [5] P. Glasserman, K. Sigman and D. Yao (Editors) (1996). Stochastic Networks: Stability and Rare Events. Springer Lecture Notes in Statistics 117, Springer- Verlag, New York.
- [6] T. Huang, and K. Sigman (1999). Unstable parameter region for FIFO Bramson-type queueing networks. (Technical Report, Columbia University, Dept. IEOR).
- [7] A. Scheller-Wolf and K. Sigman (1998). New Bounds for delay moments in multi-server queues. QUESTA, 26, 169–186.
- [8] K. Sigman (1999). (Guest Editor). Special Volume on Queues with Heavy-tailed Distribution, QUESTA, 33.
- [9] K. Sigman (2003). Sensitivity of stability to path length distributions in a FIFO multi-class network. (Technical report, Columbia University)